



# Cerone<sup>®</sup>

5L e



A growth regulator for use in winter and spring varieties of barley and winter varieties of wheat, rye and triticale

**Ethephon, (2-chloroethylphosphonic acid) is an organophosphate. Handle with care.**

## MAPP 20711

A soluble concentrate formulation containing 480 g/L (39.6% w/w) Ethephon, (2-chloroethylphosphonic acid).

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Bayer CropScience Limited  
230 Cambridge Science Park  
Milton Road, Cambridge CB4 0WB  
Telephone: 01223 226500

**For 24 hour emergency information contact Bayer CropScience Limited  
Telephone: 00800 1020 3333**

**CERONE**

Contains 480 g/l (39.6% w/w) Ethephon,  
(2-chloroethylphosphonic acid).

**Danger**

May be corrosive to metals

Causes serious eye damage

Toxic to aquatic life with long lasting effects

Wear protective gloves/protective clothing/eye protection/face protection

IF IN EYES; Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do.

Continue rinsing

Immediately call a POISON CENTRE or doctor/physician

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

**IMPORTANT INFORMATION**

FOR USE ONLY AS AN AGRICULTURAL PLANT GROWTH REGULATOR

Crops/ situations	Maximum individual dose (L product/Ha)	Maximum Total dose (L product/Ha)	Maximum total number of treatments	Latest time of application
Winter wheat	0.75	0.75	-	Before flag leaf sheath opening stage (GS 47)
Winter triticale	1.0	1.0	-	Before flag leaf sheath opening stage (GS 47)
Winter barley	1.0	1.0	-	before 1st spikelet of inflorescence just visible stage (GS 49)
Spring barley	0.5	0.5	-	before 1st spikelet of inflorescence just visible stage (GS 49)
Winter rye	1.0	1.0	-	before 1st spikelet of inflorescence just visible stage (GS 49)

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**



To access the **Safety Data Sheet** for this product scan the code or use the link below:  
<https://crops.science.bayer.co.uk/our-products/adjuvants/cerone>  
or alternatively contact your supplier

**Bayer**

## **SAFETY PRECAUTIONS**

### **OPERATOR PROTECTION**

This product contains an anticholinesterase organophosphorus compound and must not be used by those under medical advice NOT to work with such compounds

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACE SHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However engineering control may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH CONCENTRATE from skin or eyes immediately.

AVOID ALL CONTACT BY MOUTH.

WHEN USING DO NOT EAT, DRINK OR SMOKE

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

### **ENVIRONMENTAL PROTECTION**

Do not contaminate water with the product or its container. Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads.

### **STORAGE AND DISPOSAL**

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

## **DIRECTIONS FOR USE**

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Cerone plant growth regulator reduces straw length and gives greater resistance to lodging and aids harvesting of winter and spring varieties of barley and winter varieties of wheat, rye and triticale. The full benefit of Cerone treatment will only be obtained on crops which have received adequate inputs (e.g. fertilizer and fungicides).

The best response to Cerone plant growth regulator is obtained when the crop is growing vigorously. Apply Cerone plant growth regulator before lodging has started. It is recommended that only crops grown under conditions of high fertility and, therefore, at risk from lodging, should be treated with Cerone plant growth regulator. Thin crops and particularly those on soils of low fertility are less likely to lodge, and in addition such crops of winter wheat may ripen prematurely.

## RESTRICTIONS

Stress factors reduce crop growth and may also lead to secondary tillering and small grains. Cerone plant growth regulator if used under stress conditions may lead to these effects being more pronounced, particularly on spring barley.

Cereals can be susceptible to drought stress. Do not spray when the soil is very dry.

Do not use within 10 days of any herbicide or liquid fertilizer treatment.

Do not spray crops heavily diseased or suffering from pest damage, nutrient deficiency or herbicide stress.

Do not spray during cold weather or periods of night frost.

Do not spray where crop is wet or rain is imminent.

Avoid overlapping spray bouts.

Avoid drift of spray to adjoining crops.

## CROP SPECIFIC INFORMATION

Use on winter and spring varieties of barley and on winter varieties of wheat, rye and triticale.

**WINTER BARLEY:** Cerone can be applied to winter barley from the 'second node detectable' stage of crop growth (GS 32) up to and including the 'first awns just visible' stage (GS 49). The greatest resistance to lodging is achieved by treatment at the 'flag leaf just visible' stage of crop growth (GS 37).

**High Lodging Risk Crops:** For the greatest resistance to lodging use a sequential treatment of an approved formulation

containing chlormequat followed by Cerone. Apply chlormequat according to manufacturers' label instructions followed by Cerone at 0.75 L/ha from the 'flag leaf just visible' stage of crop growth (GS 37) up to and including the 'first awns just visible' stage (GS 49). The optimum time for Cerone application is at the 'flag leaf just visible' stage of crop growth (GS 37).

Where the crop has not received prior treatment with chlormequat apply Cerone at 1.0 L/ha from the 'flag leaf just visible' stage of crop growth (GS 37) up to and including 'first awns just visible' stage (GS 49).

Cerone may also be applied from the second node detectable stage of crop growth (GS 32) but the optimum time for Cerone application is at the 'flag leaf just visible' stage (GS 37).

**Low Lodging Risk Crops:** Where lodging is not expected to be a severe problem, the crop may be treated with a sequential programme of an approved formulation containing chlormequat followed by a reduced rate of Cerone. Apply chlormequat according to manufacturers' label instructions followed by Cerone at 0.5 L/ha from the 'flag leaf just visible' stage of crop growth (GS 37) up to and including 'first awns just visible' stage (GS 49).

Where the crop has not received prior treatment with chlormequat apply Cerone at 0.75 L/ha from the 'flag leaf just visible' stage of crop growth (GS 37) up to and including the 'first awns just visible' stage (GS 49). The optimum time for Cerone application is at the 'flag leaf just visible' stage of crop growth (GS 37).

**WINTER WHEAT:** For the greatest resistance to lodging use a sequential programme of an approved chlormequat product followed by Cerone. Apply chlormequat according to manufacturers label instructions, followed by Cerone at 0.5 L/ha from the 'flag leaf just visible' stage of crop growth (GS 37) up to and including the 'boots swollen' stage (GS 45). Do not spray crops where the leaf sheaths have split and the ears are visible. Where the crop has not received prior treatment with chlormequat apply Cerone at 0.75 L/ha from the 'flag leaf just visible' stage of crop growth (GS 37) up to and including the 'boots swollen' stage (GS 45). Do not spray crops where the leaf sheaths have split and the ears are visible.

**WINTER TRITICALE:** Cerone may be applied at 1.0 L/ha from the 'flag leaf just visible' stage of crop growth (GS 37) up to and including the 'boots swollen' stage (GS 45). Do not spray crops where the leaf sheaths have split and the ears are visible. The best control of lodging will be given if the crop has received prior treatment with chlormequat at the recommended timing.

**WINTER RYE:** Apply Cerone at a maximum of 1.0 L/ha from the 'flag leaf just visible' stage of crop growth (GS 37) up to and including the 'first awns just visible' stage (GS 49). Crops of winter rye which have received prior treatment with approved chlormequat products may be treated with Cerone from the 'flag leaf just visible' stage (GS 37) up to and including the 'first awns just visible' stage (GS 49).

**SPRING BARLEY:** Apply Cerone at 0.5 L/ha from the 'second node detectable' stage of crop growth (GS 32) up to and including the 'first awns just visible' stage (GS 49).

Only crops which have been drilled early, have a high yield potential and are likely to suffer severe lodging should be treated with Cerone. Crops suffering from stress or likely to be stressed, for example due to soil conditions, should not be treated (see RESTRICTIONS section above) because undesirable side effects may occur, for example secondary (green) tillers, trapped heads, or shrivelled grains. Do not use on the variety Triumph.

### **Sprayer Application**

Apply through a conventional field crop sprayer using medium nozzles at a pressure of 2-3 bar. Good spray coverage is essential. Ensure the sprayer is correctly calibrated before use and adjusted to give adequate and even coverage of the top leaves. Spray immediately after mixing. Spray quality: Cerone plant growth regulator should be applied as a MEDIUM spray (BCPC category). Water volume: 200-400 L/ha.

Cerone may also be applied in a reduced water volume of 100 litres per hectare as a FINE spray (BCPC category) provided suitable spray nozzles are selected for this type of operation.

### **MIXING**

Half fill the spray tank with clean water, add the required amount of Cerone and agitate during the completion of filling.

## **TANK CLEANING INSTRUCTIONS**

After each days work, drain the sprayer, wash thoroughly with water and liquid detergent and spray out completely.

Ensure that all liquid is removed from the sprayer tank, pump and hoses and ensure nozzles and nozzle filters are clean.

Remove nozzles, open tank and drain pump to allow free access of air to all parts of the system.

GB84406643c rA10

## CERONE

Contains 480 g/l (39.6% w/w) Ethephon, (2-chloroethylphosphonic acid).

### Danger

May be corrosive to metals

Causes serious eye damage

Toxic to aquatic life with long lasting effects

Wear protective gloves/protective clothing/eye protection/face protection

IF IN EYES; Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do.

Continue rinsing

Immediately call a POISON CENTRE or doctor/physician

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**



### IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL PLANT GROWTH REGULATOR

Crops/situations	Maximum individual dose (L product/Ha)	Maximum Total dose (L product/Ha)	Maximum total number of treatments	Latest time of application
Winter wheat	0.75	0.75	-	Before flag leaf sheath opening stage (GS 47)
Winter triticale	1.0	1.0	-	Before flag leaf sheath opening stage (GS 47)
Winter barley	1.0	1.0	-	before 1st spikelet of inflorescence just visible stage (GS 49)
Spring barley	0.5	0.5	-	before 1st spikelet of inflorescence just visible stage (GS 49)
Winter rye	1.0	1.0	-	before 1st spikelet of inflorescence just visible stage (GS 49)

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**



To access the **Safety Data Sheet** for this product scan the code or use the link below:  
<https://cropscience.bayer.co.uk/our-products/adjuvants/cerone>  
 or alternatively contact your supplier

**Bayer**