Important guidance

Sprayer best practice for optimal uptake



Emerger provides all-round pre-emergence control of broad-leaved weeds when used in accordance with the following application guidelines:



Ridges should be rounded with few clods



Emerger should be applied uniformly to the ground surface to form a homogenous film



Apply Emerger as soon as possible to settled ridges, at least 7 days before emergence of the crop



Apply via a horizontal boom sprayer, in 200-400 litres of water per hectare as a MEDIUM spray (BCPC category)



Use just one application at a maximum rate of 1.75 L/ha

Best practice for successful spray applications:

- 1 Start with a clean sprayer both inside and out
- 2 Match the nozzle/in-line filters to spray quality avoid using too fine mesh that readily blocks
- 3 Shake the container well before use
- Half fill the spray tank with clean water and begin agitation before adding the required amount of Emerger to the tank
- 5 Wash out the container and add the washings to the spray solution in the tank
- 6 Top up with remaining clean water
- 7 Maintain constant agitation at all times of any tank mix containing Emerger during filling, transportation and spraying
- 8 Do not leave the sprayer filled with the spray solution un-agitated for long periods
- **9** Always check the sprayer tank bottom between loads if sediment is seen rinse the tank and flush through the system to clear before re-filling
- Always stick to the Bayer approved Emerger tank mix sheet: cropscience.bayer.co.uk/emerger

